

	<h1>Potato Progress</h1> <p>Research &amp; Extension for the Potato Industry of Idaho, Oregon, &amp; Washington</p> <p>Andrew Jensen, Editor. <a href="mailto:ajensen@potatoes.com">ajensen@potatoes.com</a>; 509-760-4859 <a href="http://www.nwpotatoresearch.com">www.nwpotatoresearch.com</a></p>
<p>Volume XVIII, Number 9</p>	<p>16 July 2018</p>

## Vitamin B1 application on potato: an additional tool for sustainable management of Potato virus Y?

Aymeric Goyer<sup>1</sup>, Silvia I. Rondon<sup>2</sup>

<sup>1</sup> Department of Botany and Plant Pathology, Hermiston Agricultural Research and Extension Center, Oregon State University, Hermiston, OR 97838, USA

<sup>2</sup> Department of Crop and Soil Sciences, Hermiston Agricultural Research and Extension Center Oregon State University, Hermiston, OR 97838, USA

Potato has a plethora of diseases including bacteria, fungi, viruses, and nematodes. Potato productivity and production also can be compromised by arthropod pests. It is estimated that losses due to plant diseases alone can cost about \$10 billion per year (16% of potential production) [1]. Learning about how to control diseases and vectors is a major concern to the potato industry. Most control methods available to growers kill or minimize the presence and effect(s) of either the pathogen or the vector. However, there is increasing pressure to decrease the use of chemicals to reduce non-target effects, reduce impact on the environment including protecting soil health, food safety, and workers. Therefore, there is a need to search for alternative control measures. One such measure is the use of plant immunity inducers (PIIs). PIIs are immune-active compounds that can induce resistance in plants, in a similar way to vaccines in animals. There are two categories of PIIs: (1) non-biologically active molecules that are synthetic plant defense elicitors such as jasmonic acid analogs; and (2) biologically active molecules, which naturally occur in nature, mostly in plants but also in microorganisms. They include different categories of molecules such as metabolites, oligosaccharides, proteins, lipids, and microbial inducers. Some of the most recently discovered PIIs are vitamins, in particular B vitamins. There are several reports in the literature on the priming effect of, for instance, riboflavin in tobacco, rice and bean, thiamin on rice, cucumber, tobacco, soybean, and grapevine, and folic acid in *Arabidopsis* (Table 1). Thiamin also has a repellent effect on aphids in barley [2].


To our knowledge, there is no literature available that evaluates the efficiency of vitamins in potato. Thus, motivated by this lack of information, we decided to test the effect of thiamin (also known as vitamin B1) application against Potato virus Y (PVY). The study focused on evaluating the effect of thiamin concentrations and frequency of applications on disease expression.

**Table 1. Examples of reports of vitamin or vitamin precursor application to control diseases in various plants (adapted from [3]).**

<b>Vitamin</b>	<b>Plant</b>	<b>Pathogen</b>	<b>Reference</b>
Riboflavin	Tobacco	<i>Alternaria alternata</i> Keissl	[4]
		<i>Tobacco mosaic virus</i>	[5]
		<i>Phytophthora parasitica</i> L.	[5]
	Rice	<i>Rhizoctonia solani</i> Kuhn	[6]
	Bean	<i>Botrytis cinerea</i> Persoon	[7]
Thiamin	Rice	<i>Xanthomonas oryzae</i> Ishiyama	[8]
		<i>Rhizoctonia solani</i> Kuhn	[9]
	Cucumber	<i>Colletotrichum lagenarium</i> Berk. & Mont	[8]
	Tobacco	<i>Pepper mild mottle virus</i>	[8]
	Soybean	<i>Macrophomina phaseolina</i> (Tassi) Goid	[10]
	Grapevine	<i>Plasmopara viticola</i> Berl & De Toni	[11]
	Barley	Aphids	[2]
Folic acid	<i>Arabidopsis</i>	<i>Pseudomonas syringae</i> Van Hall	[12]
<i>p</i> -aminobenzoic acid	Cucumber	<i>Cucumber mosaic virus</i>	[13]
	Tobacco	<i>Pectobacterium carotovorum</i> Waldee	[14]


**What we did...**

The following controlled study was done at the Oregon State University Hermiston Agricultural Research and Extension Center in Hermiston, OR, in a greenhouse. We used disease-free Ranger Russet plantlets that we propagated in tissue culture before transferring them to soil in pots (Figure 1). Two treatments were included: A, one application of thiamin one day before PVY inoculation; B: two applications (one application one day before PVY inoculation and one 28 days after PVY inoculation). We had four different concentrations of thiamin, sprayed until run off in spray bottles with Tween 80 as surfactant. We inoculated plants with PVY N:O because this strain produces clear foliar symptoms in Ranger Russet. After inoculation, we collected leaf samples every week starting 2 weeks after PVY inoculation.



**Figure 1. Scheme showing the experimental timeline of plant growth, virus inoculation, thiamin applications, and leaf sampling.**


We sampled leaves as shown in Figure 2. Leaflet 0 was the initial inoculated leaf. We sampled the leaf just above (leaflet 1), then mid-canopy leaflets (2+3), and new emerging leaflets (4+5, and as the plants kept growing we had new emerging leaves that we also sampled 6+7, 8+9 etc...). We used ELISA to detect the virus and quantify the level of virus in the plants. We confirmed our ELISA results by using another quantification method, Real Time quantitative RT-PCR (qPCR).



**Figure 2. Diagram of leaflet sampling.**

### What we found...

Figure 3 shows the virus level in the youngest sampled leaflets based on the concentration of thiamin applied over time for one or two applications. The control was PVY-free and no thiamin treatment. All the others plant/treatments were inoculated with PVY and treated with either no thiamin or 1, 10, 50, and 100 mM thiamin. At 14 dpi, we could detect the virus when there was no thiamin or 1 mM thiamin applied, but not with higher concentrations. At 26 dpi, there was lower virus level when plants were treated with 100 mM thiamin compared to the no thiamin treatment control. This trend becomes more obvious at 35, 43, and 50 dpi. The most significant decrease in virus level was when plants were treated with two applications of thiamin at a concentration of 100 mM; in this case, the virus was not or barely detectable.



**Figure 3. Relative potato virus Y (PVY) level in leaflets sampled at various days post-inoculation (dpi) and treated with thiamin concentrations (0 mM, 1 mM, 10 mM, 50 mM, 100 mM) twice, once at the start of the experiment 24 h prior to inoculation with PVY<sup>N:O</sup>, and once 28 dpi. PVY level was determined by ELISA. All plants were treated with thiamin and inoculated with PVY<sup>N:O</sup> except the control which was not treated with thiamin and not inoculated with PVY. Data are mean ELISA values per leaflet at each sampling date for all plants where thiamin was applied twice, once at the start of the experiment 24 h prior to inoculation with PVY<sup>N:O</sup>, and once 28 dpi.**

### Conclusions

In this study, we showed that thiamin application at 100 mM significantly lowered the PVY titer throughout the plant, and two applications produced the best results. This experiment was done under controlled conditions using only one variety. Future research should determine whether thiamin application has the same effect on other commercial potato varieties and whether the economics of using this promising alternative method of control benefits the bottom line of growers' production. Our study is the first report on testing vitamins in potatoes and could serve as a model in other cropping systems. The use of this alternative way of pathogen/disease control would not replace existing control method, but would complement current practices for a more sustainable production.

This study was published in the American Journal of Potato Research [15] and was partially funded by the OSU Agricultural Research Foundation.

## References

1. Oerke EC: **Crop losses to pests.** *J Agric Sci* 2006, **144**:31-43.
2. Hamada AM, Jonsson LMV: **Thiamine treatments alleviate aphid infestations in barley and pea.** *Phytochemistry* 2013, **94**:135-141.
3. Boubakri H, Gargouri M, Mliki A, Brini F, Chong J, Jbara M: **Vitamins for enhancing plant resistance.** *Planta* 2016, **244**:529-543.
4. Zhang SJ, Yang X, Sun MW, Sun F, Deng S, Dong HS: **Riboflavin-induced Priming for Pathogen Defense in Arabidopsis thaliana.** *J Integr Plant Biol* 2009, **51**:167-174.
5. Liu F, Wei FF, Wang L, Liu H, Zhu XP, Liang YC: **Riboflavin activates defense responses in tobacco and induces resistance against Phytophthora parasitica and Ralstonia solanacearum.** *Physiol Mol Plant Pathol* 2010, **74**:330-336.
6. Taheri P, Tarighi S: **Riboflavin induces resistance in rice against Rhizoctonia solani via jasmonate-mediated priming of phenylpropanoid pathway.** *Journal of Plant Physiology* 2010, **167**:201-208.
7. Azami-Sardoei Z, Franca SC, De Vleeschauwer D, Hofte M: **Riboflavin induces resistance against Botrytis cinerea in bean, but not in tomato, by priming for a hydrogen peroxide-fueled resistance response.** *Physiol Mol Plant Pathol* 2010, **75**:23-29.
8. Ahn IP, Kim S, Lee YH: **Vitamin B1 functions as an activator of plant disease resistance.** *Plant Physiol* 2005, **138**:1505-1515.
9. Bahuguna RN, Joshi R, Shukla A, Pandey M, Kumar J: **Thiamine primed defense provides reliable alternative to systemic fungicide carbendazim against sheath blight disease in rice (Oryza sativa L.).** *Plant Physiology and Biochemistry* 2012, **57**:159-167.
10. Abdel-Monaim MF: **Role of riboflavin and thiamine in induced resistance against charcoal rot disease of soybean.** *Afr J Biotechnol* 2011, **10**:10842-10855.
11. Boubakri H, Wahab MA, Chong JL, Bertsch C, Mliki A, Soustre-Gacougnolle I: **Thiamine induced resistance to Plasmopara viticola in grapevine and elicited host-defense responses, including HR like-cell death.** *Plant Physiology and Biochemistry* 2012, **57**:120-133.
12. Wittek F, Kanawati B, Wenig M, Hoffmann T, Franz-Oberdorf K, Schwab W, Schmitt-Kopplin P, Vlot AC: **Folic acid induces salicylic acid-dependent immunity in Arabidopsis and enhances susceptibility to Alternaria brassicicola.** *Mol Plant Pathol* 2015, **16**:616-622.
13. Song GC, Choi HK, Ryu CM: **The folate precursor para-aminobenzoic acid elicits induced resistance against Cucumber mosaic virus and Xanthomonas axonopodis.** *Annals of Botany* 2013, **111**:925-934.
14. Yang SY, Park MR, Kim IS, Kim YC, Yang JW, Ryu CM: **2-Aminobenzoic acid of Bacillus sp BS107 as an ISR determinant against Pectobacterium carotovorum subsp carotovorum SCC1 in tobacco.** *Eur J Plant Pathol* 2011, **129**:371-378.
15. Vinchesi AC, Rondon SI, Goyer A: **Priming Potato with Thiamin to Control Potato Virus Y.** *Am J Potato Res* 2017, **94**:120-128.



**Potato Association  
of America**  
102nd Annual Meeting  
July 22-26, 2018  
Boise, Idaho

Visit [www.uidaho.edu/PAA2018](http://www.uidaho.edu/PAA2018)



## Special one-day registration for Industry Day, July 23<sup>rd</sup>

The Potato Association of America (PAA) will hold its 102<sup>nd</sup> annual meeting July 22-26 in Idaho's capital and the nation's top potato production state. The group is meeting in Boise to highlight the potato industry's importance in southwestern Idaho and is expected to draw more than 300 association members. "We especially hope that Idaho potato growers will turn out for the industry day," said U of I researcher Pamela Hutchinson.

Hutchinson, a weed scientist, serves on the organizing committee for the meeting with fellow CALS researcher Nora Olsen, who co-chairs the local arrangements committee with Bret Nedrow of the J.R. Simplot Co. "Monday, July 23, is Industry Day featuring presentations on topics including zebra chip, fry processing quality, and cropping systems diversity," Hutchinson said. "That afternoon will offer a symposium about the impact of quarantined pests on the potato industry." Other sessions during the meeting will focus on diseases that threaten potato quality during storage, potato handling issues that affect quality, and research to make the industry's practices more sustainable.

The group will meet at the Boise Center with other events planned at the J.R. Simplot Co.'s JUMP in downtown Boise. Participants will also enjoy whitewater rafting and celebrate on Boise's Basque Block during their visit. The meeting will offer a tour focused on Treasure Valley agriculture, including the hops and wine industry.

The Idaho Potato Commission's Big Idaho Potato Truck will be present for the event, sporting a spiffy new giant potato, Hutchinson said.

Registration information, including a one-day special for Industry Day, and more details about the meeting are available at [www.uidaho.edu/PAA2018](http://www.uidaho.edu/PAA2018).